

12

EUROPEAN PATENT APPLICATION

21 Application number: **88304577.5**

51 Int. Cl.4: **A 01 N 35/04**
/(A01N35/04,35:02)

22 Date of filing: **20.05.88**

30 Priority: **21.05.87 US 53210**

43 Date of publication of application:
23.11.88 Bulletin 88/47

64 Designated Contracting States:
BE DE ES FR GB IT NL

71 Applicant: **Surgikos, Inc.**
2500 Arbroom Boulevard P.O. Box 130
Arlington Texas 76010 (US)

72 Inventor: **Bruckner, Norman Irving**
3432 Brookshire Drive
Plano Texas 76075 (US)

Gordon, Michael David
919 Bucknell Drive
Arlington Texas 76012 (US)

Howell, Ronald Gene
3009 Kelham court
Arlington Texas 76015 (US)

74 Representative: **Jones, Alan John et al**
CARPMAELS & RANSFORD 43 Bloomsbury Square
London, WC1A 2RA (GB)

54 Disinfecting and sterilizing composition.

57 A sterilizing and disinfecting solution is disclosed. The solution contains a saturated dialdehyde such as glutaraldehyde and an aromatic dialdehyde, such as phthalaldehyde, isophthalaldehyde or terephthalaldehyde.

EP 0 292 300 A2

BEST AVAILABLE COPY

Description

DISINFECTING AND STERILIZING COMPOSITION

Field of Invention

This invention relates to improved sterilizing and disinfecting compositions which contain a saturated dialdehyde having from 2 to 6 carbon atoms and a water soluble aromatic dialdehyde. The aromatic dialdehydes include the three isomeric benzenedicarboxaldehydes: phthalaldehyde, isophthalaldehyde and terephthalaldehyde. The present composition is particularly effective against Mycobacterium tuberculosis and closely related species at 20°C.

Prior Art

Saturated dialdehyde sterilizing and disinfecting compositions are well known. Pepper et al., U.S. Patent 3,016,328; Stonehill, U.S. Patent 3,282,775; Boucher, U.S. Patent Nos. 3,708,263, 3,912,450, 3,968,248 and 3,968,250; and Buchalter, U.S. Patent No. 3,983,252 all disclose the use of glutaraldehyde in aqueous or alcoholic solutions used to disinfect or sterilize medical devices or environmental surfaces.

Winicov, U.S. Patents Nos. 4,048,336 and 4,093,744 teach sporicidal solutions which contain glutaraldehyde and a saturated monoaldehyde. Formaldehyde, acetaldehyde, propionaldehyde and butyraldehyde are specifically identified as useful monoaldehydes.

Jacobs, U.S. Patent No. 4,436,754 also disclosed sterilizing and disinfecting solutions which contain glutaraldehyde and formaldehyde.

Rein and Nolte in Zentralblatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, 1 Abt. Orig. B 168, pp. 507-516 (1979) disclose that a range of aromatic monoaldehydes and one aromatic dialdehyde, terephthalaldehyde, have bacteriostatic and fungistatic activity.

Rein, Nolte, and Zerling in Zentralblatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, 1 Abt. Orig. B 172, pp 508-519 (1981) disclose that phthalaldehyde, isophthalaldehyde and terephthalaldehyde all have bacteriostatic and fungistatic activity.

Commercially available glutaraldehyde compositions of the type disclosed in the above mentioned U.S. Patents have long been considered to be effective against a broad range of microorganisms, including Mycobacterium tuberculosis in ten (10) minutes at a temperature of 20°C. The test employed to make the determination of effectiveness was the AOAC Tuberculocidal Test, as specified in Official methods of Analysis of the Association of Official Analytical Chemists, 14th Edition, 1984, Sections 4.045-4.050. In this Test, the organism employed is Mycobacterium bovis BCG.

It is now apparent that the standard AOAC test method gives highly erratic and variable results. This test method can show that a disinfectant composition is effective against Mycobacterium bovis BCG in 10 minutes, when in fact it is much less effective than the test indicated. An improved test method, which is both reproducible and quantitative, has been developed. The new test method uses the same test organism as the above mentioned AOAC Tuberculocidal Test. In this new test method, nine milliliters (ml) of the germicide to be tested is placed in a tube, put into a water bath and allowed to come to the desired temperature. One ml of the test organism (M. bovis BCG) is added to the tube containing the germicide to be tested. At appropriate time intervals, aliquots of the germicide-cell suspension are removed and added directly to an equal volume of appropriate neutralizer and mixed thoroughly. Ten-fold dilutions of the neutralized sample are prepared with saline dilution blanks. One ml of the appropriate dilutions are collected on the surface of membrane filters having a pore size of .45 micrometers. The filters are then washed with at least 50 ml of saline. The filters are placed on agar plates and incubated in plastic bags for 15 to 20 days at 37°C. The surviving colonies are then counted. Survival curves are constructed to determine the tuberculocidal activity of the solution. The data is plotted as S/S₀ vs. time. S₀ is the initial viable count of the test organism culture and S is the viable count at each time point.

When commercial glutaraldehyde solutions are tested using the new quantitative test method, these compositions do not kill the required 1×10^5 Mycobacterium bovis BCG in 10 minutes at 20°C. The additional exposure time required for complete kill at 20°C may be as much as several hours. This exposure time becomes impractical, since the desired turn-around time for disinfection of equipment in the hospital is 30 minutes or less. In order to kill in 30 minutes a temperature of 25°C is required, and in order to kill in 10 minutes a temperature of 30°C is required.

Since the normal hospital room temperatures are between 20°C and 25°C, additional costs associated with heating conventional glutaraldehyde compositions would be required to kill all the organisms within the desired 10 to 30 minute time frame.

Summary of the Invention


It has now been discovered that the addition of relatively small amounts of an aromatic dialdehyde to the saturated dialdehyde composition provides a sterilizing and disinfecting solution which gives much more rapid kill of Mycobacterium bovis BCG at 20°C. The addition of the aromatic dialdehyde does not diminish the capability of the composition to destroy other microorganisms including the very resistant bacterial spores.

Brief Description of the Drawing

Fig. 1 is a semi-log graph plotting the ratio of the number of surviving organisms over the number of original organisms versus time for various germicidal solutions at 20°C.

Description of the Invention

The aromatic dialdehydes, which are useful in the present invention have the formula:



and are commonly called:

phthalaldehyde, where X is CHO and Y and Z are H,
isophthalaldehyde, where Y is CHO and X and Z are H, and
terephthalaldehyde, where Z is CHO and X and Y are H.

The preferred aromatic dialdehyde is phthalaldehyde because of its good solubility in water and tuberculocidal activity.

The compositions of the present invention contain a saturated dialdehyde having 2 to 6 carbon atoms as the major antimicrobial in the compositions. The saturated dialdehydes which are known to have sporicidal activity are malonaldehyde, succinaldehyde, glutaraldehyde, adipaldehyde, and oxalaldehyde. The preferred saturated dialdehyde is glutaraldehyde.

The concentration of glutaraldehyde in the composition is generally between 0.25 and 6% by weight at the dilution or concentration when the composition is used. Higher concentrations of glutaraldehyde, i.e., 10-30% may be used for shipping the composition to the point of use and the composition would then be diluted to the desired use concentration. The preferred concentration of glutaraldehyde at use is between 1% and 4%.

The aromatic dialdehyde is present in the composition, at use concentration, in amount of between 0.005% and 1% by weight based on the total weight of the composition. The preferred concentration being from 0.01% to 0.3% on the same basis. The limits on the amount of the aromatic dialdehyde used in the present composition is a function of the solubility of the particular aromatic dialdehyde in water. It is possible to increase the concentration of a particular aromatic dialdehyde in the solution by dissolving the aromatic dialdehyde in a co-solvent, miscible with water, and adding the dissolved aromatic dialdehyde to water. Suitable co-solvents include methanol, ethanol, isopropanol, polyols, tetrahydrofuran, dimethylsulfoxide and dioxane.

An alkalinizing salt is used in the composition as a buffer to maintain the proper pH of the composition during use. The alkalinizing salt may be the type disclosed in the Pepper et al., U.S. Patent No. 3,016,328, which is an alkali metal carbonate or bicarbonate, e.g., sodium bicarbonate or potassium bicarbonate, or may be a phosphate. The buffer may also be an organic carboxylate salt such as sodium citrate, sodium acetate, potassium citrate or potassium acetate. The particular salt or mixtures of salt are present in a sufficient amount, 0.1% to 2.5% based on the total weight of the solution, to give the desired pH. The pH of the composition, to obtain the optimum antimicrobial activity at room temperature, is between pH 7.0 and 9.0. However, the tuberculocidal activity enhancement provided by the aromatic dialdehyde is not dependent on pH.

The composition may contain other ingredients such as a surfactant, an odor suppressant, a corrosion inhibitor, a stabilizer for the saturated dialdehyde to inhibit polymerization, an antioxidant, sequesterant, a dye and a fragrance. The use of these other ingredients is well-known in the art.

The compositions of the present invention may be formulated in two or more parts which are combined immediately prior to use. The formulation of the composition into multiple parts extends the shelf life of the composition. The saturated dialdehydes are more effective against bacterial spores at an alkaline pH. However, the saturated dialdehydes also have a tendency to polymerize with themselves or with the aromatic dialdehyde at an alkaline pH, thereby reducing the effective concentration of both the dialdehydes in the composition. The dialdehydes of the present invention can be formulated in an aqueous solution at an acid pH, and activated with an alkalinizing agent immediately prior to use, shifting the pH to the alkaline range. This procedure is disclosed in the previously mentioned Pepper et al., U.S. Patent No. 3,016,328.

In the following Examples, all percentages are weight percentages, based on the total weight of the solutions unless otherwise indicated. The bacteriological testing was by the new tuberculocidal test methodology previously described.

Example 1

In the Example, small amounts of phthalaldehyde, isophthalaldehyde and terephthalaldehyde were added to 20% glutaraldehyde solutions and tested to show the effectiveness of the compositions in killing *Mycobacterium bovis* BCG at 20°C. The amount of isophthalaldehyde or terephthalaldehyde added was based on their solubility limit in water. The test solutions were all buffered with 0.6% dipotassium hydrogen phosphate, and the pH of the solutions was adjusted to 8 with 1N H₃PO₄. The results of these tests are shown in Table I.

Table I

10

Aromatic Dialdehyde	% Aromatic Dialdehyde (w/w)	Number of Organisms Surviving		
		0 min	10 min	20 min
None		3.8×10^5	1.3×10^4	1.2×10^3
Phthalaldehyde	0.10	2.4×10^5	0	0
Isophthalaldehyde	0.25	3.8×10^5	0	0
Terephthalaldehyde	0.10	3.8×10^5	2.0×10^1	0

25

The results indicate that the aromatic dialdehydes shown in Table I provide excellent tuberculocidal activity enhancement at 20°C.

30

Example II

In the Example, the aromatic dialdehydes were tested in aqueous solution containing no glutaraldehyde to determine the effect of the same low concentrations of aromatic dialdehydes, as in Example I, against the same organism at 20°. The solutions were buffered and the pH was adjusted to 8.0 as in Example I. The results are shown in Table II.

35

Table II

40

Aromatic Dialdehyde	% Aromatic Dialdehyde (w/w)	Number of Organisms Surviving		
		0 min.	10 min.	20 min.
Phthalaldehyde	0.10	3.8×10^5	0	0
Isophthalaldehyde	0.25	2.8×10^5	2.3×10^5	2.3×10^5
Terephthalaldehyde	0.10	2.8×10^5	3.3×10^5	4.0×10^5

50

The results show that phthalaldehyde has excellent tuberculocidal activity by itself and isophthalaldehyde and terephthalaldehyde at their water solubility limit do not have any appreciable tuberculocidal activity by themselves. Use of 20% alcohol co-solvent only modestly increase the amount of isophthalaldehyde and terephthalaldehyde in the test solutions.

Example III

A series of solutions containing 20% glutaraldehyde and various concentrations of phthalaldehyde were prepared. The solutions were buffered to pH 8 as in Example 1 and tested for their effectiveness in killing *Mycobacterium bovis* BCG at 20°C. The results are shown in Table III.

65

Table III

% Phthal- aldehyde (w/w)	Number of Organisms Surviving				5
	0 min	10 min	20 min	30 min	
0.10	2.4×10^5	0	0	0	10
0.05	2.4×10^5	2.0×10^0	0	0	
0.025	2.4×10^5	4.0×10^1	0	0	
0.01	2.4×10^5	9.4×10^2	6.0×10^0	0	15
0.005	2.4×10^5	2.1×10^3	1.3×10^2	0	
0	2.4×10^5	3.5×10^3	1.0×10^3	1.1×10^2	20

The results show that compositions containing 2% glutaraldehyde and as low as 0.005% phthalaldehyde are tuberculocidal within 30 minutes at 20°C.

Example IV

A series of solutions containing 0.75% or less of glutaraldehyde and 0.05% or less of phthalaldehyde were prepared. The solutions were buffered to pH 8 as in Example I and tested for their effectiveness in killing *Mycobacterium bovis* BCG at 20°C. The results are shown in Table IV.

Table IV

%Glutar- aldehyde (w/w)	%Phthal- aldehyde (w/w)	Number of Organisms Surviving				35
		0 min	10 min	20 min	30 min	
0.75	0.05	3.4×10^5	0	0	0	40
0.75	0.025	3.4×10^5	4.2×10^1	0	0	
0.75	0.01	3.4×10^5	2.6×10^3	7.4×10^1	0	45
0.50	0.05	3.4×10^5	0	0	0	
0.50	0.025	3.4×10^5	2.8×10^1	0	0	
0.50	0.01	3.4×10^5	3.8×10^3	3.5×10^2	1.4×10^1	50
0.25	0.05	3.4×10^5	1.2×10^2	0	0	
0.25	0.025	3.4×10^5	3.3×10^2	0	0	
0.25	0.01	3.4×10^5	5.9×10^3	1.6×10^3	2.9×10^2	55

The results show that compositions containing 0.75% glutaraldehyde and only 0.01% phthalaldehyde or 0.25% glutaraldehyde and 0.025% phthalaldehyde are tuberculocidal within 30 minutes at 20°C.

Example V

Portions of a solution containing 2% glutaraldehyde, 0.10% phthalaldehyde and 0.6% dipotassium hydrogen phosphate were adjusted to different pH levels with H_3PO_4 and KOH. The solutions were tested

against *Mycobacterium bovis* BCG at 20°C to determine the effect of pH on the effectiveness of the solutions. The results are shown in Table V.

Table V

pH	Number of Organisms Surviving			
	0 min	10 min	20 min	30 min
3	3.8×10^5	0	0	0
5	3.8×10^5	0	---	---
7	3.8×10^5	0	0	---
9	3.8×10^5	0	0	---

The results indicate that the tuberculocidal activity of compositions containing glutaraldehyde and phthalaldehyde is not pH dependent.

Example VI

A solution containing 2.0% glutaraldehyde and 0.1% phthalaldehyde was tested against a 2.0% glutaraldehyde solution to compare their effectiveness in killing spores of *Bacillus subtilis* and *Clostridium sporogenes* at 20°C in 10 hours using the standard AOAC Method (AOAC Official Methods of Analysis, 14th edition, 1984, page 72). The solutions were buffered to pH 8 as in Example I. A surfactant (Pluronic P103, BASF Wyandotte, 0.2% w/w) was added to each solution to insure wetting of the silk suture and porcelain penicylinder tests carriers. The results are shown in Table VI.

Table VI

%Glutar- aldehyde (w/w)	%Phthal- aldehyde (w/w)	Total No. of Positives (Failures/ Total No. of Tests			
		<i>B. subtilis</i>		<i>C. sporogenes</i>	
		sutures	penicylinders	sutures	penicylinders
2.0	0	0/30	0/30	0/30	0/30
2.0	0.1	0/30	0/30	0/30	0/30

The results indicate that the addition of 0.1% phthalaldehyde does not inhibit the sporicidal activity of 2% glutaraldehyde.

Fig. 1 graphically illustrates the tuberculocidal enhancement provided by the addition of small amounts of an aromatic dialdehyde to glutaraldehyde solutions. The organisms tested was *Mycobacterium bovis* BCG. The curves shown in Fig. 1 are plots of S/S_0 versus time for the following solutions (all solutions were tested at 20°C and at pH 8):

<u>Curve</u>	<u>Active Ingredients</u>	
A	2% glutaraldehyde	
B	3% glutaraldehyde	5
C	2% glutaraldehyde and 0.10% phthalaldehyde	
D	2% glutaraldehyde and 0.25% isophthalaldehyde	10
E	2% glutaraldehyde and 0.10% terephthalaldehyde	15

As indicated in Fig. 1, the presence of the aromatic dialdehydes in the solution, results in a significant reduction in the time required for total kill of the organism. 20

Claims 25

1. A sterilizing and disinfecting solution with enhanced tuberculocidal activity containing from 0.25% to 6% by weight of glutaraldehyde and from 0.005% to 1% by weight of a water soluble aromatic dialdehyde selected from the group consisting of phthalaldehyde, isophthalaldehyde and terephthalaldehyde. 30
2. The composition of claim 1 in which the aromatic dialdehyde is phthalaldehyde.
3. The composition of claim 1 in which the aromatic dialdehyde is isophthalaldehyde.
4. The composition of claim 1 in which the aromatic dialdehyde is terephthalaldehyde.
5. The composition of any one of claims 1 to 4 in which the glutaraldehyde concentration is between 1 and 4% by weight. 35
6. The composition of claim 5 in which the glutaraldehyde concentration is between 2 and 4% by weight.
7. The composition of any one of claims 1 to 6 in which the pH of the composition is in the range of pH 3 to pH 9.
8. The composition of claim 7 in which the pH of the composition is in the range of pH 7 to pH 9. 40

45


50

55

60

65

0292300



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.